



City Research Online

City, University of London Institutional Repository

Citation: Vianya-Estopa, M., Nagra, M., Cochrane, A., Retallic, N., Dunning, D., Terry, L., Lloyd, A., Wolffsohn, J. S., Alderson, A., Mouat, G., McDonnell, C., Murphy, O., Huntjens, B. ORCID: 0000-0002-4864-0723, Mayhem, M., Martin, E., Sweeney, L., Evans, K., Naroo, S., Conway, R., Simo, L., Maldonado-Codina, C., Mallon, C., Underwood, J., Trivedy, K. and Joshi, M. (2020). Optimising subjective anterior eye grading precision. Contact Lens and Anterior Eye, doi: 10.1016/j.clae.2020.03.006

This is the accepted version of the paper.

This version of the publication may differ from the final published version.

Permanent repository link: <https://openaccess.city.ac.uk/id/eprint/24191/>

Link to published version: <http://dx.doi.org/10.1016/j.clae.2020.03.006>

Copyright: City Research Online aims to make research outputs of City, University of London available to a wider audience. Copyright and Moral Rights remain with the author(s) and/or copyright holders. URLs from City Research Online may be freely distributed and linked to.

Reuse: Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

City Research Online:

<http://openaccess.city.ac.uk/>

publications@city.ac.uk

Title: Optimising Subjective Anterior Eye Grading Precision

¹Marta Vianya-Estopa PhD marta.vianya@anglia.ac.uk

²Manbir Nagra PhD manbir.nagra@port.ac.uk

³Arnold Cochrane BSc ajs.cochrane@ulster.ac.uk

⁴Neil Retallic BSc optomneil@outlook.com

⁵Dean Dunning MEd D.Dunning@bradfordcollege.ac.uk

⁶Louise Terry PhD TerryL1@cardiff.ac.uk

⁷Aoife Lloyd PhD aoife.lloyd@mckernan@dit.ie

⁸James S Wolffsohn PhD j.s.w.wolffsohn@aston.ac.uk

and members of the British & Irish University & College Contact Lens Educators (BUCCLE)

Affiliations

¹Vision and Hearing Sciences, Anglia Ruskin University, Cambridge, UK

²Optometry, University of Portsmouth, Portsmouth, UK

³School of Biomedical Sciences, Ulster University, Coleraine, UK

⁴Faculty of Biology, Medicine and Health, The University of Manchester, Manchester, UK

⁵Advanced Technology Centre, Bradford College, Bradford, UK

⁶School of Optometry and Vision Sciences, Cardiff University, Cardiff, UK

⁷School of Physics & Clinical & Optometric Sciences, Technological University Dublin, Ireland.

⁸Ophthalmic Research Group, Aston University, Birmingham, UK

Corresponding author: Prof James S Wolffsohn, Aston University, Aston Triangle, Birmingham, B4 7ET, UK j.s.w.wolffsohn@aston.ac.uk

Members of the British & Irish University & College Contact Lens Educators (BUCCLE) are: Alison Alderson and Graham Mouat (University of Bradford), Claire McDonnell and Orla Murphy (TU Dublin), Byki Huntjens and Mark Mayhew (City, University of London), Eilidh Martin (Glasgow Caledonian University), Laura Sweeney (Glasgow Caledonian University), Katherine Evans (Cardiff University), Shehzad Naroo (Aston University), Robert Conway (Anglia Ruskin University), Luisa Simo (Plymouth University), Carole Maldonado-Codina and Claire Mallon (The University of Manchester), Jo Underwood (Association of British Dispensing Opticians), Kishan Trivedy (University of Portsmouth) and Mahesh Joshi (University of Plymouth).

Abstract

Purpose: To establish the optimum grading increment which ensured parity between practitioners while maximising clinical precision.

Methods: Second year optometry students ($n=127$, 19.5 ± 1.4 years, 55% female) and qualified eye care practitioners ($n=61$, 40.2 ± 14.8 years, 52% female) had 30 seconds to grade each of bulbar, limbal and palpebral hyperaemia of the upper lid of 4 patients imaged live with a digital slit lamp under 16x magnification, diffuse illumination, with the image projected on a screen. The patients were presented in a randomised sequence 3 times in succession, during which the graders used the Efron printed grading scale once to 0.1 precision, once to 0.5 precision and once to the nearest integer grade in a randomised order. Graders were masked to their previous responses.

Results: For most grading conditions less than 20% of clinicians showed a ≤ 0.1 difference in grade from the mean. In contrast, more than 50% of the student graders and 40% of experienced graders showed a difference in grade from the mean within 0.5 for all conditions under measurement. Student precision in grading was better with both 0.1 and 0.5 grading precision than grading to the nearest unit, except for limbal hyperaemia where they performed more accurately with 0.5 unit precision grading. Limbal grading precision was not affected by grading step precision for experienced practitioners, but 0.1 and 0.5 grading precision were both better than 1.0 grading precision for bulbar hyperaemia and 0.1 grading precision was better than 0.5 grading precision and both were better than 1.0 grading precision for palpebral hyperaemia.

Conclusion: Although narrower intervals scales maximise the ability to detect smaller clinical changes, the grading increment should not exceed one standard deviation of the discrepancy between measurements. Therefore, 0.5 grading increments are recommended for subjective anterior eye physiology grading (limbal, bulbar and palpebral redness).

Keywords: grading; hyperaemia; student; eye-care practitioner; scale increments

1 Introduction

2 Since their initial introduction approximately thirty years ago, anterior eye grading scales have firmly
3 established themselves as an essential part of the eye care practitioner's (ECPs) armamentarium.
4 With usage reported at approximately 60-85% amongst ECPs [1,2] this seemingly low-tech approach
5 has had a significant impact on clinical practice. Grading scales hold several advantages over the
6 sole use of written descriptions and sketches that practitioners had previously relied upon. Grading
7 scales are quantitative, simplify the monitoring and progression of pathological and physiological
8 changes, are a universal familiar language so can be interpreted by different nationalities and across
9 health care professionals, aid in medical legal cases, and ultimately facilitate patient management.

10 While grading scales are easy to use, widely available, and considered best practice [2], they are not
11 without their limitations. Grading is subjective, associated with poor repeatability [3] and high
12 variability amongst practitioners. Grading scales are not interchangeable and the scale range varies,
13 thus grading scores will differ depending on scale used [4] with estimates reported to be higher for
14 scales which have a shorter dynamic range. [5] Further, there are concerns about the grading
15 reference images themselves. Wolffsohn [6] found grading scale images did not follow a linear
16 increase in severity, but instead followed a quadratic pattern, such that precision is greater for lower
17 severity reference images i.e. the increments between gradings are unequal. Digital versions of
18 grading scales have been produced with morphing technology [7] used to generate reference images
19 down to 0.1 scale grade increments, but any improvement in grading variability has not been
20 published.

21 Some of the shortcomings may be attributable to the process of grading itself; typically, anterior eye
22 grading involves the application of a discrete scale (a limited fixed number of grades) to a continuous
23 variable (the severity of a particular ocular condition). [8] Several sources [2,8] have advocated the
24 reduction of grading scale increment size to increase clinical precision i.e. grading to the nearest
25 integer should produce poorer clinical precision than grading to the nearest 0.5 or 0.1.
26 Nevertheless, achieving adequate clinical precision may not necessitate use of the smallest grading
27 increment possible. Peterson and Wolffsohn [3] showed a mean difference of approximately 0.70-
28 1.03 bulbar redness (Efron) image grades was needed for it to be discernible by subjective grading.
29 Given the widespread use of grading scales, and their vulnerability to subjective bias, it is of clinical
30 interest to establish an evidence base for a best practice approach to grading. The aim of this study
31 was to establish the optimum grading increment which ensured parity between practitioners while
32 maximising clinical precision. Based on previously published data, it is hypothesised that whole
33 integer grading will be less accurate (a larger absolute deviation from the mean practitioner grade)
34 than grading to the nearest 0.5 or 0.1 unit.

35 Method

The study was granted a favourable ethical opinion by Ulster University (practitioner study) and Aston University (student study) ethics committees and followed the tenets of the Declaration of Helsinki. Participants gave written informed consent to take part after an explanation of the study.

The graders were 2nd year undergraduate optometry students enrolled at Aston University (n=127, 19.5 ± 1.4 years, 55% female) and qualified eye care practitioners (at least 2 years) attending the BCLA UK conference in June 2018 (n=61, 40.2 ±14.8 years, 52% female) all familiar with using grading scales with the Efron grading scale. Data collection for the two cohorts occurred on separate occasions.

The ocular surface of 4 patients with no ocular pathology were observed live under 16x magnification, diffuse illumination, with a digital slit-lamp (Keeler, Windsor, UK) and the image projected on a screen. The patients were presented in a randomised sequence 3 times in succession during which the graders used the Efron printed grading scale once to 0.1 increments, once to 0.5 increments and once to the nearest integer grade in randomised order. They had 30 seconds to grade each of bulbar, limbal and upper lid palpebral hyperaemia each time, and were masked to their previous grades.

Statistical Analysis

The absolute average difference from the mean of all graders, for each grader with each increment level was calculated for each of the 4 patients. As the data was not normally distributed, non-parametric statistics were applied (Friedman test repeated measure analysis of variance with Wilcoxon signed-rank test post-hoc pairwise comparison where significance was identified). In addition the discrepancies between pairs of observers were assessed for each of the 4 patients and the standard deviation calculated.

Based on a standard deviation of 0.4 [9] for subjective grading, a clinically significant difference (p<0.05) of 0.2 units between groups could be detected with 80% power with a sample size of 61 participants in each group and 0.15 units with 127 participants in each group.

<https://www.stat.ubc.ca/~rollin/stats/ssize/n2.html>

Results

Across the 4 patients examined, the average bulbar grade ranged from 0.8-1.5, average limbal grade ranged from 0.4 to 1.2 and the average palpebral grade ranged from 0.4 to 1.6 and was similar between patients used for the student grading and practitioner grading sessions. The distribution of the difference from the mean is shown in Figure 1 for student graders and Figure 2 for qualified eye care practitioners. The mean of these differences for each feature is shown in Table 1, along with statistical significance. There was a significant difference ($p < 0.001$) across all grade increment comparisons except practitioner graded limbal hyperaemia ($p = 0.478$). The percentage of clinicians increased for all conditions within a greater difference in grade from the mean. For most conditions less than 20% of clinicians showed a ≤ 0.1 difference in grade from the mean. In contrast, more than 50% of the student graders and 40% of experienced graders showed a difference in grade from the mean within 0.5 for all conditions under measurement.

Grading							
Increment		0.1	↔	0.5	↔	1.0	↔ 0.1
		mean	p	mean	p	Mean	P
Student n=127	Bulbar	0.40±0.30	0.342	0.42±0.31	<0.001	0.51±0.29	<0.001
	Limbal	0.44±0.31	0.156	0.42±0.31	0.001	0.47±0.36	0.259
	Palpebral	0.35±0.26	0.645	0.34±0.32	<0.001	0.49±0.27	<0.001
Practitioner n=61	Bulbar	0.58±0.50	0.633	0.58±0.53	0.004	0.64±0.45	0.001
	Limbal	0.54±0.46	0.790	0.54±0.49	0.940	0.53±0.52	0.874
	Palpebral	0.71±0.64	0.026	0.75±0.67	<0.001	0.82±0.64	<0.001

Table 1: Mean grade difference (\pm S.D.) from mean and significance between grading increments. The arrows above the significance (p) values point to the two increments being compared.

Student precision in grading was better with both 0.1 and 0.5 grading increments than grading to the nearest unit, except for limbal hyperaemia where it was only better with 0.5 unit increment grading (there was no significant difference between the 0.1 and 1.0 increments for this feature). Limbal grading precision was not affected by grading step increment for experienced practitioners, but 0.1 and 0.5 grading increments were both better than the 1.0 grading increment for bulbar hyperaemia. For palpebral hyperaemia, the 0.1 grading increment was more accurate than the 0.5 grading increment and both were better than 1.0 grading increment (Table 1). The standard deviation of discrepancies between observers was 0.65-0.87 across the students and was 0.72 to 0.84 across experienced practitioners.

Figure 1: What proportion of student clinicians were within 0.1 to 1.0 grades different from the mean of all clinicians for bulbar, limbal, and palpebral hyperaemia with each of the grading increments. N=127.

Figure 2: What proportion of experienced practitioners within 0.1 to 1.0 grades different from the mean of all clinicians for bulbar, limbal and palpebral hyperaemia with each of the grading increments. N=61.

Discussion

This study set out to show that smaller grading increment steps would lead to more accurate grading compared to the mean. In practical terms, the grades recorded by a practitioner should be as close as possible to the mean of other practitioners (average difference) rather than the discrepancy analysis (the difference between 2 practitioners) as modelled by Bailey et al. [8]. However, while this was the case for 0.5 grading units compared to whole integer grading, this was generally not the case for 0.1 grading units compared to 0.5. As shown in Table 1, the average difference from the mean was around 0.30 for student graders and 0.55 for experienced graders. The standard deviation between random pairs of observers was higher, as expected, being 0.72 for student graders and 0.78 for experienced graders. Bailey et al. [8] suggested that if the scale increment exceeds the standard deviation of the discrepancy this will result in a sharp broadening of the confidence limits. Thus, these findings suggest that a 0.5 grading step might be as precise as is possible to get when evaluating hyperaemia in the anterior eye using the Efron printed grading scale.

It is worth noting that limbal hyperaemia grading was more variable in grade than bulbar and palpebral redness. This finding is not surprising as the exact extent of the limbal region is not clearly defined clinically and graders might have been influenced by nearby conjunctival redness. Yet, observers need to ensure enough attention is given to this structure given the response between limbal hyperaemia and contact lens wear. For instance, several studies have shown that hydrogel lens wear results in significantly greater levels of limbal hyperaemia compared to silicone hydrogel lens wear for both daily and extended wear modalities, whereas bulbar redness is not significantly affected. [10-13]

Efron et al [4] suggested that grading of contact lens complications would be expected to improve with experience. His group also found grading variability improves statistically (but not clinically significant) with some experience, but no added benefit could be derived from supplemental training [14]. However, this study found experienced practitioners were less accurate than second year undergraduate optometry students. Similar findings between students and experienced practitioners were also noted by Wolffsohn et al [4]. Although a priori one might expect experienced

practitioners to show greater precision than students, this might no longer be the case as the importance of grading in the assessment of anterior eye is currently emphasised to undergraduate optometry students. Similarly, Cardona and Serés [15] noted that contact lens knowledge improved grading precision in optometry students. The students taking part in this study had received a 1 hour seminar on the principals behind grading and had used the Efron grading scales in 5 weekly 2 hour clinics. Differences in the data projection of the images, such as screen resolution and ambient brightness could have made a difference between cohorts, but the student graders used the same conditions as half of the experienced graders and the difference between them was still evident. Future work should further explore the relationship that knowledge, training and experience might have on the uses of grading scales in anterior eye and contact lens assessment. A survey of UK practitioners in 2015 [2] indicated that 91.6% of respondents used grading scales for bulbar conjunctival hyperaemia and 77.8% and 63.4% for limbal and palpebral hyperaemia respectively. It could be hypothesised that less familiarity of usage might lead to more variability with grading and this seems to be the case with practitioners.

Recently, alternative methods to subjective assessment of bulbar and limbal hyperaemia have been proposed using software such as Keratograph 5M (Oculus) that objectively detects hyperaemia. [16] Artificial intelligence learning algorithms have been applied to retinal images, demonstrating their ability not just to quantify disease changes, but also to identify other features that might differentiate disease and its progression such as tortuosity, pallor and blood flow, not traditionally utilised by clinicians. [17] However, technological advances are not yet readily available by most clinicians. In addition, the results of this new technology might not be interchangeable with results obtained using subjective grading scales. [18-19] Thus, it is important to continue to support clinicians using grading scales optimally, although, digital photography can allow direct comparison at subsequent visits and is preferable to grading.

It is important to note that this study was conducted using projected slit lamp videos of eyes without pathology. The patients examined were different between the students and experienced practitioners, but the average grades were similar for each of the ocular anterior eye features examined and the comparison was the individual's difference from the mean, so the actual mean should not have a significant effect on the results. The mean grade of each feature was ≤ 2 for each participant; the entire range of the grading scale used was not included in the study. Therefore the conclusions cannot be extended to grading precision for more severe hyperaemic cases.

In conclusion, this study showed that 0.5 grading increments should be recommended when assessing anterior eye grading (limbal, bulbar and palpebral hyperaemia). This contradicts previous recommendation by Efron et al [4] and Wolffsohn et al. [2] of recording clinical signs using 0.1 increments between grades. Although narrower intervals scales maximise the ability to detect smaller clinical changes, Bailey et al [8] also indicated that for moderate precision the grading

159 increment should not exceed one standard deviation of the discrepancy between measurements.
160 Although narrower increments have been recommended in clinical practice, Efron et al [4] and
161 Wolffsohn et al [2] found graders tended to grade using whole and half-digits indicating a reluctance
162 to use finer increments. Thus, this research provides the evidence for clinicians to adopt 0.5
163 increments in their clinical grading alongside previous research highlighting the importance of
164 recording the scale used and having the scale present when grading. [2,6]

165

166

References

- [1] Efron N, Pritchard N, Brandon K, Copeland J, Godfrey R, Hamlyn B, Vrbancic V. A survey of the use of grading scales for contact lens complications in optometric practice. *Clin Exp Optom* 2011;94:193-9.
- [2] Wolffsohn JS, Naroo SA, Christie C, Morris J, Conway R, Maldonado-Codina C. Anterior eye health recording. *Contact Lens Ant Eye* 2015;38:266-71.
- [3] Peterson RC, Wolffsohn JS. Sensitivity and reliability of objective image analysis compared to subjective grading of bulbar hyperaemia. *Br J Ophthalmol* 2007;91:1464-6.
- [4] Efron N, Morgan PB, Katsara SS. Validation of grading scales for contact lens complications. *Ophthalmic Physiol Opt.* 2001;21:17-29.
- [5] Schulze MM, Hutchings N, Simpson TL. Grading bulbar redness using cross-calibrated clinical grading scales. *Invest Ophthalmol Vis Sci* 2011;52:5812-7.
- [6] Wolffsohn JS. Incremental nature of anterior eye grading scales determined by objective image analysis. *Br J Ophthalmol* 2004; 88:1434-8.
- [7] Efron N, Morgan PB, Jagpal R. Validation of computer morphs for grading contact lens complications. *Ophthalmic Physiol Opt* 2002;22:341-49.
- [8] Bailey IL, Bullimore MA, Raasch TW and Taylor HR. Clinical Grading and the Effects of Scaling. *Invest Ophthalmol Vis Sci* 1991;32:422–32.
- [9] Efron N. Grading scales for contact lens complications. *Ophthalmic Physiol Opt* 1998;18:182-6.
- [10] Brennan NA, Coles ML, Connor HR, McIlroy RG. A 12-month prospective clinical trial of comfilcon A silicone-hydrogel contact lenses worn on a 30-day continuous wear basis. *Cont Lens Anterior Eye* 2007;30:108-18.
- [11] Dumbleton K, Keir N, Moezzi A, Feng Y, Jones L, Fonn D. Objective and subjective responses in patients refitted to daily-wear silicone hydrogel contact lenses. *Optom Vis Sci* 2006; 83:758-68.
- [12] Maldonado-Codina C, Morgan PB, Schnider CM, Efron N. Short-term physiologic response in neophyte subjects fitted with hydrogel and silicone hydrogel contact lenses. *Optom Vis Sci* 2004; 81:911-21.
- [13] Dillehay SM, Miller MB. Performance of Lotrafilcon B silicone hydrogel contact lenses in experienced low-Dk/t daily lens wearers. *Eye Contact Lens* 2007; 33:272-77.

- 196 [14] Efron N, Morgan PB, Farmer C, Furuborg J, Struk R, Carney LG. Experience and training as
 197 determinants of grading reliability when assessing the severity of contact lens complications.
 198 Ophthalmic Physiol Opt 2003;23:119-24.
- 199 [15] Cardona G & Serés C. Grading Contact Lens Complications: The Effect of Knowledge on Grading
 200 Accuracy. Curr Eye Res 2009; 34: 1074–81.
- 201 [16] Wu S, Hong J, Tian L, Cui X, Sun X and Xu J. Assessment of Bulbar Redness with a Newly
 202 Developed Keratograph. Optom Vis Sci 2015; 92: 892-899.
- 203 [17] Varadarajan AV, Poplin R, Blumer K, Angermueller CA, Ledsam J, Chopra R, Keane PA, Corrado
 204 GS, Peng L and Webster DR. Deep learning for predicting refractive error from retinal fundus images.
 205 Invest Ophthalmol Vis Sci 2018;59: 2861-8.
- 206 [18] Perez-Bartolome F and Garcia-Feijoo J. Assessment of ocular redness measurements obtained
 207 with keratography 5M and correlation with subjective grading scales. Journal Français
 208 d’Ophthalmologie 2018; 41 (9): 836-46.
- 209 [19] Huntjens B, Basi M, Nagra M. Evaluating a new objective grading software for conjunctival
 210 hyperaemia. 2019. Contact Lens Ant Eye (in press)

211

212 **Acknowledgements**

213 BUCCLE’s mission is to enhance optometry education in the UK and in this pursuit is funded by
 214 industry including Alcon, Bausch and Lomb, Coopervision, No7 Contact lenses and David
 215 Thomas/Menicon. This work was also supported by Cheryl Donnelly and the BCLA. Other BUCCLE
 216 members include Alison Alderson and Graham Mouat (University of Bradford), Claire McDonnell and
 217 Orla Murphy (TU Dublin), Byki Huntjens (City, University of London), Mark Mayhem (City, University
 218 of London), Eilidh Martin (Glasgow Caledonian University), Laura Sweeney (Glasgow Caledonian
 219 University), Katherine Evans (Cardiff University), Shehzad Naroo (Aston University), Robert Conway
 220 (Anglia Ruskin University), Luisa Simo (Plymouth University), Carole Maldonado-Codina and Claire
 221 Mallon (The University of Manchester), Jo Underwood (Association of British Dispensing Opticians),
 222 Kishan Trivedy (University of Portsmouth) and Mahesh Joshi (University of Plymouth).